



2

Description of Motion in One Dimension



- Preeti reached the metro station and found that the escalator was not working. She walked up the stationary escalator in time t_1 . On other days, if she remains stationary on the moving escalator, then the escalator takes her up in time t_2 . The time taken by her to walk up on the moving escalator will be

(a) $\frac{t_1 t_2}{t_2 - t_1}$ (b) $\frac{t_1 t_2}{t_2 + t_1}$
 (c) $t_1 - t_2$ (d) $\frac{t_1 + t_2}{2}$
(NEET 2017)
- Two cars P and Q start from a point at the same time in a straight line and their positions are represented by $x_p(t) = (at + bt^2)$ and $x_q(t) = (ft - t^2)$. At what time do the cars have the same velocity?

(a) $\frac{a-f}{1+b}$ (b) $\frac{a+f}{2(b-1)}$
 (c) $\frac{a+f}{2(1+b)}$ (d) $\frac{f-a}{2(1+b)}$
(NEET-II 2016)
- If the velocity of a particle is $v = At + Bt^2$, where A and B are constants, then the distance travelled by it between 1 s and 2 s is

(a) $\frac{3}{2}A + \frac{7}{3}B$ (b) $\frac{A}{2} + \frac{B}{3}$
 (c) $\frac{3}{2}A + 4B$ (d) $3A + 7B$
(NEET-I 2016)
- A particle of unit mass undergoes one-dimensional motion such that its velocity varies according to $v(x) = \beta x^{-2n}$, where β and n are constants and x is the position of the particle. The acceleration of the particle as a function of x , is given by

(a) $-2\beta^2 x^{-2n-1}$ (b) $-2n\beta^2 e^{-4n-1}$
 (c) $-2n\beta^2 x^{-2n-1}$ (d) $-2n\beta^2 x^{-4n-1}$
(2015 Cancelled)
- A stone falls freely under gravity. It covers distances h_1 , h_2 and h_3 in the first 5 seconds, the next 5 seconds and the next 5 seconds respectively. The relation between h_1 , h_2 and h_3 is

(a) $h_2 = 3h_1$ and $h_3 = 3h_2$
 (b) $h_1 = h_2 = h_3$
 (c) $h_1 = 2h_2 = 3h_3$
 (d) $h_1 = \frac{h_2}{3} = \frac{h_3}{5}$ *(NEET 2013)*
- The displacement 'x' (in meter) of a particle of mass 'm' (in kg) moving in one dimension under the action of a force, is related to time 't' (in sec) by $t = \sqrt{x} + 3$. The displacement of the particle when its velocity is zero, will be

(a) 4 m (b) 0 m (zero)
 (c) 6 m (d) 2 m
(Karnataka NEET 2013)
- The motion of a particle along a straight line is described by equation $x = 8 + 12t - t^3$ where x is in metre and t in second. The retardation of the particle when its velocity becomes zero is

(a) 24 m s^{-2} (b) zero
 (c) 6 m s^{-2} (d) 12 m s^{-2} (2012)
- A boy standing at the top of a tower of 20 m height drops a stone. Assuming $g = 10 \text{ m s}^{-2}$, the velocity with which it hits the ground is

(a) 10.0 m/s (b) 20.0 m/s
 (c) 40.0 m/s (d) 5.0 m/s (2011)
- A particle covers half of its total distance with speed v_1 and the rest half distance with speed v_2 . Its average speed during the complete journey is

(a) $\frac{v_1 + v_2}{2}$ (b) $\frac{2v_1 v_2}{v_1 + v_2}$
 (c) $\frac{2v_1 v_2}{v_1 + v_2}$ (d) $\frac{v_1^2 v_2^2}{v_1^2 + v_2^2}$
(Mains 2011)




10. A particle moves a distance x in time t according to equation $x = (t + 5)^{-1}$. The acceleration of particle is proportional to
 (a) (velocity)^{3/2} (b) (distance)²
 (c) (distance)⁻² (d) (velocity)^{2/3} (2010)
11. A ball is dropped from a high rise platform at $t = 0$ starting from rest. After 6 seconds another ball is thrown downwards from the same platform with a speed v . The two balls meet at $t = 18$ s. What is the value of v ?
 (Take $g = 10 \text{ m/s}^2$)
 (a) 75 m/s (b) 55 m/s
 (c) 40 m/s (d) 60 m/s (2010)
12. A particle starts its motion from rest under the action of a constant force. If the distance covered in first 10 seconds is S_1 and that covered in the first 20 seconds is S_2 , then
 (a) $S_2 = 3S_1$ (b) $S_2 = 4S_1$
 (c) $S_2 = S_1$ (d) $S_2 = 2S_1$ (2009)
13. A bus is moving with a speed of 10 ms^{-1} on a straight road. A scooterist wishes to overtake the bus in 100 s. If the bus is at a distance of 1 km from the scooterist, with what speed should the scooterist chase the bus?
 (a) 40 m s^{-1} (b) 25 m s^{-1}
 (c) 10 m s^{-1} (d) 20 m s^{-1} (2009)
14. A particle moves in a straight line with a constant acceleration. It changes its velocity from 10 ms^{-1} to 20 ms^{-1} while passing through a distance 135 m in t second. The value of t is
 (a) 12 (b) 9
 (c) 10 (d) 1.8 (2008)
15. The distance travelled by a particle starting from rest and moving with an acceleration $\frac{4}{3} \text{ m s}^{-2}$, in the third second is
 (a) $\frac{10}{3} \text{ m}$ (b) $\frac{19}{3} \text{ m}$
 (c) 6m (d) 4m (2008)
16. A particle moving along x -axis has acceleration f , at time t , given by $f = f_0 \left(1 - \frac{t}{T}\right)$, where f_0 and T are constants. The particle at $t = 0$ has zero velocity. In the time interval between $t = 0$ and the instant when $f = 0$, the particle's velocity (v_s) is
 (a) $\frac{1}{2} f_0 T^2$ (b) $f_0 T^2$
 (c) $\frac{1}{2} f_0 T$ (d) $f_0 T$ (2007)
17. A car moves from X to Y with a uniform speed v_u and returns to Y with a uniform speed v_d . The average speed for this round trip is
 (a) $\sqrt{v_u v_d}$ (b) $\frac{v_d v_u}{v_d + v_u}$
 (c) $\frac{v_u + v_d}{2}$ (d) $\frac{2v_d v_u}{v_d + v_u}$ (2007)
18. The position x of a particle with respect to time t along x -axis is given by $x = 9t^2 - t^3$ where x is in metres and t in seconds. What will be the position of this particle when it achieves maximum speed along the $+x$ direction?
 (a) 54 m (b) 81 m
 (c) 24 m (d) 32 m. (2007)
19. Two bodies A (of mass 1 kg) and B (of mass 3 kg) are dropped from heights of 16 m and 25 m, respectively. The ratio of the time taken by them to reach the ground is
 (a) 4/5 (b) 5/4
 (c) 12/5 (d) 5/12. (2006)
20. A car runs at a constant speed on a circular track of radius 100 m, taking 62.8 seconds for every circular lap. The average velocity and average speed for each circular lap respectively is
 (a) 10 m/s, 0 (b) 0, 0
 (c) 0, 10 m/s (d) 10 m/s, 10 m/s. (2006)
21. A particle moves along a straight line OX . At a time t (in seconds) the distance x (in metres) of the particle from O is given by $x = 40 + 12t - t^3$. How long would the particle travel before coming to rest?
 (a) 16 m (b) 24 m
 (c) 40 m (d) 56 m. (2006)
22. A ball is thrown vertically upward. It has a speed of 10 m/sec when it has reached one half of its maximum height. How high does the ball rise?
 (Take $g = 10 \text{ m/s}^2$)
 (a) 10 m (b) 5 m
 (c) 15 m (d) 20 m. (2005)



23. The displacement x of a particle varies with time t as $x = ae^{-\alpha t} + be^{\beta t}$, where a, b, α and β are positive constants. The velocity of the particle will
- be independent of β
 - drop to zero when $\alpha = \beta$
 - go on decreasing with time
 - go on increasing with time. (2005)
24. A man throws balls with the same speed vertically upwards one after the other at an interval of 2 seconds. What should be the speed of the throw so that more than two balls are in the sky at any time? (Given $g = 9.8 \text{ m/s}^2$)
- more than 19.6 m/s
 - at least 9.8 m/s
 - any speed less than 19.6 m/s
 - only with speed 19.6 m/s. (2003)
25. If a ball is thrown vertically upwards with speed u , the distance covered during the last t seconds of its ascent is
- ut
 - $\frac{1}{2}gt^2$
 - $ut - \frac{1}{2}gt^2$
 - $(u + gt)t$. (2003)
26. A particle is thrown vertically upward. Its velocity at half of the height is 10 m/s, then the maximum height attained by it ($g = 10 \text{ m/s}^2$)
- 8 m
 - 20 m
 - 10 m
 - 16 m. (2001)
27. Motion of a particle is given by equation $s = (3t^3 + 7t^2 + 14t + 8) \text{ m}$. The value of acceleration of the particle at $t = 1 \text{ sec}$ is
- 10 m/s²
 - 32 m/s²
 - 23 m/s²
 - 16 m/s². (2000)
28. A car moving with a speed of 40 km/h can be stopped by applying brakes after at least 2 m. If the same car is moving with a speed of 80 km/h, what is the minimum stopping distance?
- 4 m
 - 6 m
 - 8 m
 - 2 m. (1998)
29. A rubber ball is dropped from a height of 5 m on a plane. On bouncing it rises to 1.8 m. The ball loses its velocity on bouncing by a factor of
- $\frac{3}{5}$
 - $\frac{2}{5}$
 - $\frac{16}{25}$
 - $\frac{9}{25}$. (1998)
30. The position x of a particle varies with time, (t) as $x = at^2 - bt^3$. The acceleration will be zero at time t is equal to
- $\frac{a}{3b}$
 - zero
 - $\frac{2a}{3b}$
 - $\frac{a}{b}$. (1997)
31. If a car at rest accelerates uniformly to a speed of 144 km/h in 20 sec. it covers a distance of
- 1440 cm
 - 2980 cm
 - 20 m
 - 400 m. (1997)
32. A body dropped from a height h with initial velocity zero, strikes the ground with a velocity 3 m/s. Another body of same mass dropped from the same height h with an initial velocity of 4 m/s. The final velocity of second mass, with which it strikes the ground is
- 5 m/s
 - 12 m/s
 - 3 m/s
 - 4 m/s. (1996)
33. The acceleration of a particle is increasing linearly with time t as bt . The particle starts from origin with an initial velocity v_0 . The distance travelled by the particle in time t will be
- $v_0t + \frac{1}{3}bt^2$
 - $v_0t + \frac{1}{2}bt^2$
 - $v_0t + \frac{1}{6}bt^3$
 - $v_0t + \frac{1}{3}bt^3$. (1995)
34. The water drop falls at regular intervals from a tap 5 m above the ground. The third drop is leaving the tap at instant the first drop touches the ground. How far above the ground is the second drop at that instant?
- 3.75 m
 - 4.00 m
 - 1.25 m
 - 2.50 m. (1995)
35. A car accelerates from rest at a constant rate α for some time after which it decelerates at a constant rate β and comes to rest. If total time elapsed is t , then maximum velocity acquired by car will be
- $\frac{(\alpha^2 - \beta^2)t}{\alpha\beta}$
 - $\frac{(\alpha^2 + \beta^2)t}{\alpha\beta}$
 - $\frac{(\alpha + \beta)t}{\alpha\beta}$
 - $\frac{\alpha\beta t}{\alpha + \beta}$. (1994)
36. A particle moves along a straight line such that its displacement at any time t is given by $s = (t^3 - 6t^2 + 3t + 4) \text{ metres}$. The velocity when the acceleration is zero is
- 3 m/s
 - 42 m/s
 - 9 m/s
 - 15 m/s. (1994)




37. The velocity of train increases uniformly from 20 km/h to 60 km/h in 4 hours. The distance travelled by the train during this period is
 (a) 160 km (b) 180 km
 (c) 100 km (d) 120 km. (1994)
38. The displacement-time graph of a moving particle is shown below. The instantaneous velocity of the particle is negative at the point



- (a) E (b) F (c) C (d) D
 (1994)

39. A body starts from rest, what is the ratio of the distance travelled by the during the 4th and 3rd second?
 (a) $\frac{7}{5}$ (b) $\frac{5}{7}$
 (c) $\frac{7}{3}$ (d) $\frac{3}{7}$. (1993)

40. Which of the following curve does not represent motion in one dimension?



- (1992)
41. A body dropped from top of a tower fall through 40 m during the last two seconds of

- its fall. The height of tower is ($g = 10 \text{ m/s}^2$)
 (a) 60 m (b) 45 m
 (c) 80 m (d) 50 m.

(1992)

42. A car moves a distance of 200 m. It covers the first half of the distance at speed 40 km/h and the second half of distance at speed v . The average speed is 48 km/h. The value of v is
 (a) 56 km/h (b) 60 km/h
 (c) 50 km/h (d) 48 km/h.

(1991)

43. A bus travelling the first one-third distance at a speed of 10 km/h, the next one-third at 20 km/h and at last one-third at 60 km/h. The average speed of the bus is
 (a) 9 km/h (b) 16 km/h
 (c) 18 km/h (d) 48 km/h.

(1991)

44. A car covers the first half of the distance between two places at 40 km/h and another half at 60 km/h. The average speed of the car is
 (a) 40 km/h (b) 48 km/h
 (c) 50 km/h (d) 60 km/h.

(1990)

45. What will be the ratio of the distance moved by a freely falling body from rest in 4th and 5th seconds of journey?
 (a) 4 : 5 (b) 7 : 9
 (c) 16 : 25 (d) 1 : 1.

(1989)

46. A car is moving along a straight road with a uniform acceleration. It passes through two points P and Q separated by a distance with velocity 30 km/h and 40 km/h respectively. The velocity of the car midway between P and Q is
 (a) 33.3 km/h (b) $20\sqrt{2}$ km/h
 (c) $25\sqrt{2}$ km/h (d) 35 km/h.

(1988)